

A really simple elementary proof of the uniform boundedness theorem

Alan D. Sokal*
Department of Physics
New York University
4 Washington Place
New York, NY 10003 USA
sokal@nyu.edu

May 7, 2010

revised October 20, 2010

to appear in the *American Mathematical Monthly*

Abstract

I give a proof of the uniform boundedness theorem that is elementary (i.e., does not use any version of the Baire category theorem) and also extremely simple.

Key Words: Uniform boundedness; gliding hump; sliding hump; Baire category.

Mathematics Subject Classification (MSC 2000) codes: 46B99 (Primary); 46B20, 46B28 (Secondary).

*Also at Department of Mathematics, University College London, London WC1E 6BT, England.

One of the pillars of functional analysis is the uniform boundedness theorem:

UNIFORM BOUNDEDNESS THEOREM. Let \mathcal{F} be a family of bounded linear operators from a Banach space X to a normed linear space Y . If \mathcal{F} is pointwise bounded (i.e., $\sup_{T \in \mathcal{F}} \|Tx\| < \infty$ for all $x \in X$), then \mathcal{F} is norm-bounded (i.e., $\sup_{T \in \mathcal{F}} \|T\| < \infty$).

The standard textbook proof (e.g., [17, p. 81]), which goes back to Stefan Banach, Hugo Steinhaus, and Stanisław Saks in 1927 [3], employs the Baire category theorem or some variant thereof.¹ This proof is very simple, but its reliance on the Baire category theorem makes it not completely elementary.

By contrast, the original proofs given by Hans Hahn [7] and Stefan Banach [2] in 1922 were quite different: they began from the assumption that $\sup_{T \in \mathcal{F}} \|T\| = \infty$ and used a “gliding hump” (also called “sliding hump”) technique to construct a sequence (T_n) in \mathcal{F} and a point $x \in X$ such that $\lim_{n \rightarrow \infty} \|T_n x\| = \infty$.² Variants of this proof were later given by T. H. Hildebrandt [11] and Felix Hausdorff [9, 10].³ These proofs are elementary, but the details are a bit fiddly.

Here is a *really* simple proof along similar lines:

Lemma. Let T be a bounded linear operator from a normed linear space X to a normed linear space Y . Then for any $x \in X$ and $r > 0$, we have

$$\sup_{x' \in B(x, r)} \|Tx'\| \geq \|T\|r, \quad (1)$$

where $B(x, r) = \{x' \in X : \|x' - x\| < r\}$.

PROOF. For $\xi \in X$ we have

$$\max\{\|T(x + \xi)\|, \|T(x - \xi)\|\} \geq \frac{1}{2}[\|T(x + \xi)\| + \|T(x - \xi)\|] \geq \|T\xi\|, \quad (2)$$

where the second \geq uses the triangle inequality in the form $\|\alpha - \beta\| \leq \|\alpha\| + \|\beta\|$. Now take the supremum over $\xi \in B(0, r)$. \square

PROOF OF THE UNIFORM BOUNDEDNESS THEOREM. Suppose that $\sup_{T \in \mathcal{F}} \|T\| = \infty$, and choose $(T_n)_{n=1}^\infty$ in \mathcal{F} such that $\|T_n\| \geq 4^n$. Then set $x_0 = 0$, and for $n \geq 1$ use the lemma to choose inductively $x_n \in X$ such that $\|x_n - x_{n-1}\| \leq 3^{-n}$ and $\|T_n x_n\| \geq \frac{2}{3}3^{-n}\|T_n\|$. The sequence (x_n) is Cauchy, hence convergent to some $x \in X$; and it is easy to see that $\|x - x_n\| \leq \frac{1}{2}3^{-n}$ and hence $\|T_n x\| \geq \frac{1}{6}3^{-n}\|T_n\| \geq \frac{1}{6}(4/3)^n \rightarrow \infty$. \square

¹ See [4, p. 319, note 67] concerning credit to Saks.

² Hahn’s proof is discussed in at least two modern textbooks: see [14, Exercise 1.76, p. 49] and [13, Exercise 3.15, pp. 71–72].

³ See also [18, pp. 63–64] and [21, pp. 74–75] for an elementary proof that is closely related to the standard “nested ball” proof of the Baire category theorem; and see [8, Problem 27, pp. 14–15 and 184] and [12] for elementary proofs in the special case of linear functionals on a Hilbert space.

Remarks. 1. As just seen, this proof is most conveniently expressed in terms of a *sequence* (x_n) that converges to x . This contrasts with the earlier “gliding hump” proofs, which used a *series* that sums to x . Of course, sequences and series are equivalent, so each proof can be expressed in either language; it is a question of taste which formulation one finds simpler.

2. This proof is extremely wasteful from a quantitative point of view. A quantitatively sharp version of the uniform boundedness theorem follows from Ball’s “plank theorem” [1]: namely, if $\sum_{n=1}^{\infty} \|T_n\|^{-1} < \infty$, then there exists $x \in X$ such that $\lim_{n \rightarrow \infty} \|T_n x\| = \infty$ (see also [15]).

3. A similar (but slightly more complicated) elementary proof of the uniform boundedness theorem can be found in [6, p. 83].

4. “Gliding hump” proofs continue to be useful in functional analysis: see [20] for a detailed survey.

5. The standard Baire category method yields a slightly stronger version of the uniform boundedness theorem than the one stated here, namely: if $\sup_{T \in \mathcal{F}} \|Tx\| < \infty$ for a *nonmeager* (i.e., second category) set of $x \in X$, then \mathcal{F} is norm-bounded.

6. The uniform boundedness theorem has generalizations to suitable classes of non-normable and even non-metrizable topological vector spaces (see, e.g., [19, pp. 82–87]). I leave it to others to determine whether any ideas from this proof can be carried over to these more general settings.

7. More information on the history of the uniform boundedness theorem can be found in [4, pp. 302, 319n67], [5, pp. 138–142], and [16, pp. 21–22, 40–43].

Acknowledgments. I wish to thank Keith Ball for reminding me of Hahn’s “gliding hump” proof; it was my attempt to fill in the details of Keith’s sketch that led to the proof reported here. Keith informs me that versions of this proof have been independently devised by at least four or five people, though to his knowledge none of them have bothered to publish it. I also wish to thank David Edmunds and Bob Megginson for helpful correspondence, and three anonymous referees for valuable suggestions concerning the exposition. Finally, I wish to thank Jürgen Voigt and Markus Haase for drawing my attention to the elementary proofs in [6, 18, 21].

This research was supported in part by U.S. National Science Foundation grant PHY-0424082.

References

- [1] K. Ball, The plank problem for symmetric bodies, *Invent. Math.* **104** (1991) 535–543.
- [2] S. Banach, Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales, *Fund. Math.* **3** (1922) 133–181; also in *Œuvres avec des Commentaires*, vol. 2, Éditions scientifiques de Pologne, Warsaw, 1979.

- [3] S. Banach and H. Steinhaus, Sur le principe de la condensation des singularités, *Fund. Math.* **9** (1927) 50–61.
- [4] G. Birkhoff and E. Kreyszig, The establishment of functional analysis, *Historia Math.* **11** (1984) 258–321.
- [5] J. Dieudonné, *History of Functional Analysis*, North-Holland, Amsterdam, 1981.
- [6] I. Gohberg, S. Goldberg, and M. A. Kaashoek, *Basic Classes of Linear Operators*, Birkhäuser, Basel, 2003.
- [7] H. Hahn, Über Folgen linearer Operationen, *Monatsh. Math. Phys.* **32** (1922) 3–88.
- [8] P. R. Halmos, *A Hilbert Space Problem Book*, 2nd ed., Springer-Verlag, New York, 1982.
- [9] F. Hausdorff, Zur Theorie der linearen metrischen Räume, *J. Reine Angew. Math.* **167** (1932) 294–311.
- [10] J. Hennefeld, A nontopological proof of the uniform boundedness theorem, *Amer. Math. Monthly* **87** (1980) 217.
- [11] T. H. Hildebrandt, On uniform limitedness of sets of functional operations, *Bull. Amer. Math. Soc.* **29** (1923) 309–315.
- [12] S. S. Holland Jr., A Hilbert space proof of the Banach-Steinhaus theorem, *Amer. Math. Monthly* **76** (1969) 40–41.
- [13] B. D. MacCluer, *Elementary Functional Analysis*, Springer-Verlag, New York, 2009.
- [14] R. E. Megginson, *An Introduction to Banach Space Theory*, Springer-Verlag, New York, 1998.
- [15] V. Müller and J. Vršovský, Orbits of linear operators tending to infinity, *Rocky Mountain J. Math.* **39** (2009) 219–230.
- [16] A. Pietsch, *History of Banach Spaces and Linear Operators*, Birkhäuser, Boston, 2007.
- [17] M. Reed and B. Simon, *Methods of Modern Mathematical Physics*, vol. I, *Functional Analysis*, Academic Press, New York, 1972.
- [18] F. Riesz and B. Sz.-Nagy, *Functional Analysis*, Ungar, New York, 1955.
- [19] H. H. Schaefer and M. P. Wolff, *Topological Vector Spaces*, 2nd ed., Springer-Verlag, New York, 1999.

- [20] C. Swartz, *Infinite Matrices and the Gliding Hump*, World Scientific, River Edge, NJ, 1996.
- [21] J. Weidmann, *Linear Operators in Hilbert Spaces*, Springer-Verlag, New York, 1980.